



# Parent/Athlete Concussion Information Sheet

A concussion is a type of traumatic brain injury that changes the way the brain normally works. A concussion is caused by bump, blow, or jolt to the head or body that causes the head and brain to move rapidly back and forth. Even a “ding,” “getting your bell rung,” or what seems to be a mild bump or blow to the head can be serious.

## WHAT ARE THE SIGNS AND SYMPTOMS OF CONCUSSION?

Signs and symptoms of concussion can show up right after the injury or may not appear or be noticed until days or weeks after the injury.

If an athlete reports **one or more** symptoms of concussion listed below after a bump, blow, or jolt to

### Did You Know?

- Most concussions occur *without* loss of consciousness.
- Athletes who have, at any point in their lives, had a concussion have an increased risk for another concussion.
- Young children and teens are more likely to get a concussion and take longer to recover than adults.

the head or body, s/he should be kept out of play the day of the injury and until a health care professional, experienced in evaluating for concussion, says s/he is symptom-free and it’s OK to return to play.

SIGNS OBSERVED BY COACHING STAFF	SYMPTOMS REPORTED BY ATHLETES
Appears dazed or stunned	Headache or “pressure” in head
Is confused about assignment or position	Nausea or vomiting
Forgets an instruction	Balance problems or dizziness
Is unsure of game, score, or opponent	Double or blurry vision
Moves clumsily	Sensitivity to light
Answers questions slowly	Sensitivity to noise
Loses consciousness ( <i>even briefly</i> )	Feeling sluggish, hazy, foggy, or groggy
Shows mood, behavior, or personality changes	Concentration or memory problems
Can’t recall events <i>prior</i> to hit or fall	Confusion
Can’t recall events <i>after</i> hit or fall	Just not “feeling right” or “feeling down”



## CONCUSSION DANGER SIGNS

In rare cases, a dangerous blood clot may form on the brain in a person with a concussion and crowd the brain against the skull. An athlete should receive immediate medical attention if after a bump, blow, or jolt to the head or body s/he exhibits any of the following danger signs:

- One pupil larger than the other
- Is drowsy or cannot be awakened
- A headache that not only does not diminish, but gets worse
- Weakness, numbness, or decreased coordination
- Repeated vomiting or nausea
- Slurred speech
- Convulsions or seizures
- Cannot recognize people or places
- Becomes increasingly confused, restless, or agitated
- Has unusual behavior
- Loses consciousness (*even a brief loss of consciousness should be taken seriously*)

## WHY SHOULD AN ATHLETE REPORT THEIR SYMPTOMS?

If an athlete has a concussion, his/her brain needs time to heal. While an athlete's brain is still healing, s/he is much more likely to have another concussion. Repeat concussions can increase the time it takes to recover. In rare cases, repeat concussions in young athletes can result in brain swelling or permanent damage to their brain. *They can even be fatal.*

It's better to miss one game than the whole season. For more information on concussions, visit: [www.cdc.gov/Concussion](http://www.cdc.gov/Concussion).

### *Remember*

Concussions affect people differently. While most athletes with a concussion recover quickly and fully, some will have symptoms that last for days, or even weeks. A more serious concussion can last for months or longer.

## WHAT SHOULD YOU DO IF YOU THINK YOUR ATHLETE HAS A CONCUSSION?

If you suspect that an athlete has a concussion, remove the athlete from play and seek medical attention. Do not try to judge the severity of the injury yourself. Keep the athlete out of play the day of the injury and until a health care professional, experienced in evaluating for concussion, says s/he is symptom-free and it's OK to return to play.

Rest is key to helping an athlete recover from a concussion. Exercising or activities that involve a lot of concentration, such as studying, working on the computer, or playing video games, may cause concussion symptoms to reappear or get worse. After a concussion, returning to sports and school is a gradual process that should be carefully managed and monitored by a health care professional.

\_\_\_\_\_  
Student-Athlete Name Printed

\_\_\_\_\_  
Student-Athlete Signature

\_\_\_\_\_  
Date

\_\_\_\_\_  
Parent or Legal Guardian Printed

\_\_\_\_\_  
Parent or Legal Guardian Signature

\_\_\_\_\_  
Date



# Keep Their Heart in the Game

## A Sudden Cardiac Arrest Information Sheet for Athletes and Parents/Guardians

### What is sudden cardiac arrest?

Sudden cardiac arrest (SCA) is when the heart stops beating, suddenly and unexpectedly. When this happens blood stops flowing to the brain and other vital organs. SCA is NOT a heart attack. A heart attack is caused by a blockage that stops the flow of blood to the heart. SCA is a malfunction in the heart's electrical system, causing the victim to collapse. The malfunction is caused by a congenital or genetic defect in the heart's structure.

### How common is sudden cardiac arrest in the United States?

As the leading cause of death in the U.S., there are more than 300,000 cardiac arrests outside hospitals each year, with nine out of 10 resulting in death. Thousands of sudden cardiac arrests occur among youth, as it is the #2 cause of death under 25 and the #1 killer of student athletes during exercise.

### Who is at risk for sudden cardiac arrest?

SCA is more likely to occur during exercise or physical activity, so student-athletes are at greater risk. While a heart condition may have no warning signs, studies show that many young people do have symptoms but neglect to tell an adult. This may be because they are embarrassed, they do not want to jeopardize their playing time, they mistakenly think they're out of shape and need to train harder, or they simply ignore the symptoms, assuming they will "just go away." Additionally, some health history factors increase the risk of SCA.

**FAINTING**  
is the  
**#1 SYMPTOM**  
OF A HEART CONDITION

### What should you do if your student-athlete is experiencing any of these symptoms?

We need to let student-athletes know that if they experience any SCA-related symptoms it is crucial to alert an adult and get follow-up care as soon as possible with a primary care physician. If the athlete has any of the SCA risk factors, these should also be discussed with a doctor to determine if further testing is needed. Wait for your doctor's feedback before returning to play, and alert your coach, trainer and school nurse about any diagnosed conditions.

### What is an AED?

An automated external defibrillator (AED) is the only way to save a sudden cardiac arrest victim. An AED is a portable, user-friendly device that automatically diagnoses potentially life-threatening heart rhythms and delivers an electric shock to restore normal rhythm. Anyone can operate an AED, regardless of training. Simple audio direction instructs the rescuer when to press a button to deliver the shock, while other AEDs provide an automatic shock if a fatal heart rhythm is detected. A rescuer cannot accidentally hurt a victim with an AED—quick action can only help. AEDs are designed to only shock victims whose hearts need to be restored to a healthy rhythm. Check with your school for locations of on-campus AEDs.



## The Cardiac Chain of Survival

On average it takes EMS teams up to 12 minutes to arrive to a cardiac emergency. Every minute delay in attending to a sudden cardiac arrest victim decreases the chance of survival by 10%. Everyone should be prepared to take action in the first minutes of collapse.

### Early Recognition of Sudden Cardiac Arrest



Collapsed and unresponsive.  
Gasping, gurgling, snorting, moaning  
or labored breathing noises.  
Seizure-like activity.

### Early Access to 9-1-1



Confirm unresponsiveness.  
Call 9-1-1 and follow emergency  
dispatcher's instructions.  
Call any on-site Emergency Responders.

### Early CPR



Begin cardiopulmonary resuscitation  
(CPR) immediately. Hands-only CPR involves fast  
and continual two-inch chest compressions—  
about 100 per minute.

### Early Defibrillation



Immediately retrieve and use an automated  
external defibrillator (AED) as soon as possible  
to restore the heart to its normal rhythm. Mobile  
AED units have step-by-step instructions for a by-  
stander to use in an emergency situation.

### Early Advanced Care



Emergency Medical Services (EMS)  
Responders begin advanced life support  
including additional resuscitative measures and  
transfer to a hospital.



# Keep Their Heart in the Game

## Recognize the Warning Signs & Risk Factors of Sudden Cardiac Arrest (SCA)

Tell Your Coach and Consult Your Doctor if These Conditions are Present in Your Student-Athlete

### Potential Indicators That SCA May Occur

- Fainting or seizure, especially during or right after exercise
- Fainting repeatedly or with excitement or startle
- Excessive shortness of breath during exercise
- Racing or fluttering heart palpitations or irregular heartbeat
- Repeated dizziness or lightheadedness
- Chest pain or discomfort with exercise
- Excessive, unexpected fatigue during or after exercise

### Factors That Increase the Risk of SCA

- Family history of known heart abnormalities or sudden death before age 50
- Specific family history of Long QT Syndrome, Brugada Syndrome, Hypertrophic Cardiomyopathy, or Arrhythmogenic Right Ventricular Dysplasia (ARVD)
- Family members with unexplained fainting, seizures, drowning or near drowning or car accidents
- Known structural heart abnormality, repaired or unrepaired
- Use of drugs, such as cocaine, inhalants, "recreational" drugs, excessive energy drinks or performance-enhancing supplements

### What is CIF doing to help protect student-athletes?

CIF amended its bylaws to include language that adds SCA training to coach certification and practice and game protocol that empowers coaches to remove from play a student-athlete who exhibits fainting—the number one warning sign of a potential heart condition. A student-athlete who has been removed from play after displaying signs or symptoms associated with SCA may not return to play until he or she is evaluated and cleared by a licensed health care provider. Parents, guardians and caregivers are urged to dialogue with student-athletes about their heart health and everyone associated with high school sports should be familiar with the cardiac chain of survival so they are prepared in the event of a cardiac emergency.

I have reviewed and understand the symptoms and warning signs of SCA and the new CIF protocol to incorporate SCA prevention strategies into my student's sports program.

STUDENT-ATHLETE SIGNATURE

PRINT STUDENT-ATHLETE'S NAME

DATE

PARENT/GUARDIAN SIGNATURE

PRINT PARENT/GUARDIAN'S NAME

DATE

For more information about Sudden Cardiac Arrest visit

California Interscholastic Federation  
<http://www.cifstate.org>

Eric Paredes Save A Life Foundation  
<http://www.epsavealife.org>

CardiacWise (20-minute training video)  
<http://www.sportsafetyinternational.org>

