






Administering Glucagon

Call 911 immediately upon administration



1. Position the student on his or/ her side.
2. Remove the cap from the glass vial.
3. Pull the needle cover off the syringe.
4. Insert the needle into the vial and inject liquid.
5. Swirl to dissolve.
6. Withdraw the glucagon solution back into the syringe and remove the needle from the vial.
7. Check for air bubbles in the syringe. Tap any visible air to the top of the syringe and gently push on the plunger until the air is removed.
8. Insert the needle at a 90-degree angle and inject the glucagon into a large muscle (upper arm, thigh, or upper outer area of the buttock).
9. Withdraw the needle and apply slight pressure to the injection site.
10. Keep the student positioned on his or her side.
- 11.
12. Remain with the student until Emergency Medical Services (EMS) assumes control.

BAQSIMI GLUCAGON ADMINISTRATION



STEPS OF ADMINISTRATION

1. Verify that the student is unconscious or unable to swallow.
2. Take Baqsimi out of the container.
3. Hold device between fingers and thumb. **DO NOT** push plunger yet.



4. Insert tip gently into one nostril until fingers touch the outside of the nose.



5. Push plunger firmly all the way in. Dose is complete when the green line disappears.
6. Call 911 or have someone else call 911 immediately.
7. Stay with student until EMS arrives.
8. **Document Baqsimi glucagon administration using CCSD Medication Administration Log.**

GVOKE GLUCAGON ADMINISTRATION



1. Verify that the student is unconscious or unable to swallow.
2. Call 911 or have someone else call 911 immediately.
3. Tear open pouch at dotted line.
4. Remove syringe
5. Choose injection site and expose skin. Lower abdomen, outer thighs and outer upper arm are all appropriate sites.
6. Pull needle off cap. **DO NOT REMOVE AIR BUBBLES!**
7. Pinch the skin and insert the needle at a 90° angle.
8. Push plunger down to inject glucagon.
9. Turn patient on side

Gvoke HypoPen (glucagon injection) 1 mg Auto-Injector

1. Prepare → **2. Inject** → **3. Assist**

1. Prepare
Tear Open Pouch at Dotted Line. Remove Auto-Injector.
Pull off Red Cap.

2. Inject
Choose Injection Site and Expose Skin.
Front View Back View
Lower Abdomen, Outer Thigh, or Outer Upper Arm
Push Down on Skin to Start. Hold Down for 5 Seconds.
Wait for Window to Turn Red.
Hold Down for 5 Sec.

3. Assist
Turn Patient on Side. Call Emergency Medical Help.

After the Injection, Put the Used Pen in a Safe Place Until It Can Be Disposed of Into a FDA Cleared Sharps Container.

Red Cap **Needle End**

10. Stay with the student until EMS arrives.
11. Document Gvoke glucagon administration using CCSD Medication Administration Log.

ZEGALOUGE AUTOINJECTOR (Glucagon) ADMINISTRATION INSTRUCTIONS

Storage information

- Store ZEGALOUGE in the refrigerator between 36°F to 46°F (2°C to 8°C).
- **Do not** freeze.
- Store ZEGALOUGE in the red protective case it comes in.
- ZEGALOUGE can also be stored at room temperature between 68°F to 77°F (20°C to 25°C) for up to 12 months. See example to the right for how to keep track of this 12-month period.
- **Do not** return to refrigerator after storing at room temperature.
- Throw away ZEGALOUGE if it has been stored at room temperature for more than 12 months.
- Replace ZEGALOUGE before the expiration date printed on the red protective case.
- **Keep ZEGALOUGE and all medicines out of the reach of children.**

If ZEGALOUGE is removed from the refrigerator to be stored at room temperature, write down the new 12-month expiration date on the label found on the red protective case.



For example: If ZEGALOUGE was removed from refrigerator to be stored at room temperature in January 2020, then write the new expiration date of January 2021 on the label.

Began storing at room temperature

JAN	FEB	MAR	APR	MAY	JUN	JUL
	1	2	3	4	5	6
AUG	SEP	OCT	NOV	DEC	JAN	
7	8	9	10	11	12	

Expires in 12 months

The image shows the ZEGALOUGE autoinjector label with the new expiration date JAN 2021 written in the field. A yellow box highlights the new date, and another yellow box highlights the current date SEP2021. Below the label is a yellow box with the text 'Throw away on earlier date'.

Note: ZEGALOUGE will expire on the earlier of the two expiration dates.

Description of parts


Red protective case




ZEGALOGUE

Autoinjector

Before injection




After injection



Before injection

Choose injection site and expose skin




Injection sites include:

- **Outer upper arms**
- **Lower abdomen**
(at least 2 inches from the belly button)
- **Front or back of thighs**
- **Buttocks**

Roll back any clothing to expose bare skin. **Do not** inject through clothes.

Hold the red protective case upright and remove gray lid




- Hold the red protective case upright with the gray lid on top.
- Pull the gray lid up to open.
- Carefully remove ZEGALOGUE from the red protective case without dropping it.

How to inject

Step 1

Remove the gray cap from needle end



Pull the gray cap straight off.


Do not put your hand or fingers anywhere near the yellow needle guard. Touching the yellow needle guard may cause an accidental needle stick injury.

Step 2

Push and hold down ZEGALOGUE for 10 seconds and check window is red



Push ZEGALOGUE straight down on skin until the yellow needle guard is fully pressed down. You may hear the first click.




Keep holding ZEGALOGUE down and slowly count to 10 seconds. During this time the **medicine window will turn red** and you may hear a second click.

Check that the medicine window is red, which means that the full dose has been given.

Step 3

Remove ZEGALOGUE from injection site



Remove ZEGALOGUE by lifting it straight up.

The yellow needle guard will cover the needle and lock, preventing an accidental needle stick injury.

After injection

- After you have given the injection, roll the unconscious person on to their side to prevent choking.
- **Call for emergency medical help or a healthcare provider right away after you have injected ZEGALOGUE.** Even if the ZEGALOGUE injection helps the person to wake up, still call for emergency medical help right away. If the person does not respond after 15 minutes, another dose may be given, if available.
- Once the person is able to safely consume food or drink, give the person a fast-acting source of sugar (such as fruit juice) and a long-acting source of sugar (such as crackers with cheese or peanut butter).
- **Replace the used ZEGALOGUE right away so you will have a new ZEGALOGUE in case you need it.**

Hypoglycemia may happen again after receiving ZEGALOGUE treatment.

Early symptoms of hypoglycemia may include:

- sweating
- drowsiness
- dizziness
- sleep disturbances
- irregular heartbeat (palpitation)
- anxiety
- tremor
- blurred vision
- hunger
- slurred speech
- restlessness
- depressed mood
- tingling in the hands, feet, lips, or tongue
- irritability
- abnormal behavior
- lightheadedness
- unsteady movement
- inability to concentrate
- personality changes
- headache

ZEGALOGUE PREFILLED SYRINGE (Glucagon) ADMINISTRATION INSTRUCTIONS


Storage information

- Store ZEGALOGUE in the refrigerator between 36°F to 46°F (2°C to 8°C).
- **Do not** freeze.
- Store ZEGALOGUE in the red protective case it comes in.
- ZEGALOGUE can also be stored at room temperature between 68°F to 77°F (20°C to 25°C) for up to 12 months. See example to the right for how to keep track of this 12-month period.
- **Do not** return to refrigerator after storing at room temperature.
- Throw away ZEGALOGUE if it has been stored at room temperature for more than 12 months.
- Replace ZEGALOGUE before the expiration date printed on the red protective case.
- **Keep ZEGALOGUE and all medicines out of the reach of children.**

If ZEGALOGUE is removed from the refrigerator to be stored at room temperature, write down the new 12-month expiration date on the label found on the red protective case.



For example: If ZEGALOGUE was removed from refrigerator to be stored at room temperature in January 2020, then write the new expiration date of January 2021 on the label.



Note: ZEGALOGUE will expire on the earlier of the two expiration dates.

Description of parts

Red protective case




ZEGALOGUE[®]

Prefilled Syringe



Before injection

Choose injection site and expose skin




Injection sites include:

- **Outer upper arms**
- **Lower abdomen**
(at least 2 inches from the belly button)
- **Front or back of thighs**
- **Buttocks**

Roll back any clothing to expose bare skin. **Do not** inject through clothes.

Hold the red protective case upright and remove gray cap




- Hold the red protective case upright with the gray cap on top.
- Pull the gray cap up to open.
- Carefully remove ZEGALOGUE from the red protective case without dropping it.

How to inject

Step 1


Remove the gray needle cover



Pull the gray needle cover straight off. Be careful not to bend the needle.

Step 2


Pinch the skin and insert the needle



Gently pinch the skin and insert the entire needle into the skin at a 45° angle.

Step 3


Give the injection



After inserting the needle, release the pinched skin and slowly press the plunger rod all the way down until the syringe is empty and the plunger rod stops.

Step 4

Remove the needle



After the plunger rod stops and the injection is complete, carefully remove the needle from the injection site.